



Is there a relationship between temporomandibular disorder and sacroiliac joint dysfunction?

Is TMD associated with SJD?

Sultan İğrek^{1,2}, Tuğba Kuru Çolak³, İlker Çolak⁴, Erdinç Çekiç⁵

¹ Department of Physiotherapy and Rehabilitation, Institute of Health Sciences, Marmara University, İstanbul, Türkiye.

² Department of Therapy and Rehabilitation, Vocational School, İstanbul Beykent University, İstanbul, Türkiye.

³ Department of Physiotherapy and Rehabilitation, Faculty of Health Sciences, Marmara University, İstanbul, Türkiye.

⁴ Clinic of Orthopaedics and Traumatology, VM Medical Park Maltepe Hospital, İstanbul, Türkiye.

⁵ Otorhinolaryngology Private Clinic, İstanbul, Türkiye.

Abstract

Aim: The aim of this prospective clinical study was to evaluate the possible association between temporomandibular disorders (TMD) and sacroiliac joint dysfunction (SIJD) and to determine the immediate effect of a single session of orofacial manual therapy (OMT) on SIJD.

Methods: Twenty-four patients, aged 18 to 60, diagnosed with both TMD and SIJD, underwent a single-session OMT treatment. Orofacial pain, pressure pain thresholds (PPT) in the masseter, temporalis, and spina iliaca posterior superior (SIPS), temporomandibular joint (TMJ) range of motion (ROM), hipomobility in the SIPS, and functionality were evaluated before and after the treatment.

Results: A statistically significant improve was observed in the Orofacial pain scores, all TMJ ROM values, PPT in masseter, temporalis, and SIPS, Patrick's Faber test score, and Oswestry Disability Index score after treatment in all participants ($p < 0.001$).

Conclusion: Assessing and managing the joints together can lead to a more comprehensive evaluation of patients with Sacroiliac and Temporomandibular joint dysfunction symptoms, enhancing the overall effectiveness of the treatment.

Keywords

temporomandibular disorder, sacroiliac joint, facial pain, manual therapies

DOI: 10.4328/ACAM.22373 Ann Clin Anal Med 2025;16(2):99-104

Received: 19.08.2024

Accepted: 21.10.2024

Published Online: 06.11.2024

Printed: 01.02. 2025

Corresponding Author: Sultan İğrek, Department of Therapy and Rehabilitation, Vocational School, İstanbul Beykent University, İstanbul, Türkiye.

E-mail: sultanigrek@gmail.com P: +90 537 210 06 54

Corresponding Author ORCID ID: <https://orcid.org/0000-0002-6983-0470>

Other Authors ORCID ID: Tuğba Kuru Çolak, <https://orcid.org/0000-0002-3263-2278> . İlker Çolak, <https://orcid.org/0000-0003-2960-2825>

Erdinç Çekiç, <https://orcid.org/0000-0001-8784-174X>

Introduction

Temporomandibular disorder (TMD) is a specific category of orofacial disorders that encompasses issues such as temporomandibular joint (TMJ) pain, fatigue in the craniocervical muscles—particularly the masticatory muscles—restricted movement of the jaw, and clicking or deviation during jaw movements.¹ TMD is a prevalent issue in clinical settings, affecting approximately 33% of the general population.² The most frequently reported symptom is facial pain, which can present as either acute or chronic.¹ The causes of TMD are diverse and include factors such as anxiety, stress, bruxism, alterations in occlusion and posture, dysfunction of the masticatory muscles and other bodily structures, or a combination of these elements.³

Sacroiliac joint dysfunction (SIJD) is one of the most common causes of non-discogenic low back pain. With a reported incidence of 10-27%, it is also responsible for mechanical and chronic hip or leg pain.⁴ The relationship between occlusal disorders of the TMJ, cervical spine and sacroiliac joint (SIJ) has been evaluated and coexisting dysfunction in these regions has been obtained.⁵ Cases of TMD have been reported in which lumbosacral pain and malocclusion were successfully treated together.^{6,7} These results suggest a functional connection between these regions. This connection is based on diaphragms, neural connections, and the presence of myofascial chains.^{8,9} Based on these findings, this study focused on the possible relationship between TMD and SIJD. The aim of this prospective clinical study was to evaluate the possible association between TMD and SIJD, and to determine the immediate effect of a single session of orofacial manual therapy (OMT) on SIJD. The hypothesis was that a single session of manual therapy applied to the temporomandibular joint can reduce complaints related to the sacroiliac joint.

Materials and Methods

Study Design and Participants

This study was conducted on patients aged 18–60 years diagnosed with TMD - reduction disc displacement between September 2023 and March 2024. This study was approved by Ethical Committee. All the study procedures conformed to the provisions of the World Medical Association Declaration of Helsinki. Patients who agreed to participate in the study were included in the study if they gave verbal and written informed consent. This study was registered in the WHO International Clinical Trials Registry Platform (NCT06035341).

Patients with orofacial pain who presented to the Department of Otorhinolaryngology at Outpatient Clinic and were diagnosed with Temporomandibular Disorder with reduction disc displacement, based on physical and radiological examinations conducted by a specialist physician, were subsequently referred to the Orthopedics and Traumatology Department. In this department, the patients were evaluated for sacroiliac joint dysfunction (SIJD) by an Orthopedics and Traumatology specialist. Those patients who were diagnosed with SIJD in addition to Temporomandibular Disorder with reduction disc displacement and who agreed to participate in the study were included in the research.

Patients were excluded from the study if they had a history of malignant condition, trauma or surgery in the cranial or cervical region, are uncooperative, use regular analgesic or anti-inflammatory drugs, have dentofacial anomalies, active inflammatory arthritis, metabolic diseases (Gout, osteoporosis, Cushing’s disease, and hyper/hypo parathyroidism), connective tissue, rheumatological (Systemic lupus erythematosus and

scleroderma) and hematological disorders (Anemia and leukemia), diagnosed psychiatric diseases, and received TMDrelated physical therapy less than 6 months ago.

Assessment Methods

Demographic and clinical characteristics of patients were recorded. As evaluation parameters orofacial pain, pressure pain threshold (PPT) in masseter, temporalis, and spina iliaca posterior superior (SIPS), TMJ range of motion (ROM), functionality and mobility in SIJ were evaluated. Orofacial pain, PPT, TMJ ROM and mobility in SIJ were evaluated before treatment and 5 minutes after treatment. Functionality was evaluated before treatment and 1 week after treatment. Orofacial pain was evaluated using the Visual Analog Scale (VAS). This scale consists of numbers from 0 to 10. Accordingly, “0” describes no pain, and “10” describes unbearable pain. The patient was asked to mark the intensity of the pain he/she felt and it was explained that the intensity of pain increased from 0 to 10.¹⁰

Mandibular movements (depression, right and left lateral excursion, protrusion, and retrusion) using a 15 cm ruler. Individuals were asked to perform these movements as much as possible without causing pain. The distance between the maxillary and mandibular central incisors were measured using a ruler. Measurements were recorded three times and the average of the recorded measurements were noted.¹¹

A digital algometer (Jtech Medical Commander Algometer) was used to objectively measure the PPT. The algometer was checked by pressing the pulp of the thumb of the hand before starting the measurement. Then, pressure was applied 1 cm below the SIPS and on the central TP of the masseter and temporalis muscles by increasing the pressure by 1 kg/ cm² every three seconds until the individual feels pain. The physiotherapist passively supported the individual’s head with the other hand. The individual was instructed to report when they felt pain while applying force with the device (kg/cm²). Three pressure pain threshold measurements were made




Figure 1. Orofacial manual therapy




Figure 2. Flow chart

with a 2-min rest interval between trials. The mean of the 3 trials was calculated and used for analysis.¹²

The Patrick-Faber test was used to assess the mobility of the sacroiliac joint. For the measurement of the test, one examiner stabilized the patient while another measured the vertical distance in centimeters between the mark on the lateral edge of the patella and the base. A greater distance indicated higher tension in the adductor muscles and less sacroiliac joint mobility.⁵

Functionality was assessed using the Oswestry Disability Index (ODI). The ODI is a valid and reliable questionnaire that evaluates daily living activities across 10 different areas, including pain intensity, personal care, lifting, walking, sitting, standing, sleep, changes in pain, social life, and travel.¹³ Scores range from a minimum of 0 to a maximum of 50. The questionnaire was filled out in person prior to treatment and was administered via telephone one week post-treatment.

Treatment Protocol

Participants underwent a single session of OMT (Figure 1). Immediately after the treatment, the participants were asked to walk for 5 minutes and then orofacial pain, pressure pain threshold, TMJ ROM and SIJ mobility were assessed as a second evaluation. Functionality was assessed 1 week later as the third evaluation. The OMT treatment program included trigger point therapy, myofascial release, TMJ traction, joint mobilization with traction, muscle energy technique, mandibular fascia release, occipital release, and ligament therapy.¹⁴⁻¹⁵

Ethical Approval

This study was approved by the Ethics Committee of Marmara University (Date: 05.05.2023, Decision No: 09.2023.675).

Statistical Analysis

G*Power V3.1.9.2 was used to calculate the sample size of the study. The effect size calculated based on the study of Fink et al. (Faber test pre-intervention = 16.3 ± 3.8, postintervention = 18.4 ± 3.8) was determined as f = 0.55.⁵ Accordingly, the minimum sample size to was reached with a power of 0.80 for α = 0.05 type 1 error probability was calculated as 23 people.

Table 1. Sociodemographic and clinical data of the study participants

Parameter		X ± SD
Age, y		34.8 ± 13.1
Weight, kg		70.3 ± 16.1
BMI (kg/m²)		26.2 ± 5.5
Height		1.63 ± 0.1
Parameter		X ± SD
Sex	Female	21 (87.5%)
	Male	3 (12.5%)
Education level	Education level below high school	13 (54.2%)
	High school and above education level	11 (45.8%)
Trauma history	Yes	10 (41.7%)
	No	14 (58.3%)
Surgery history	Yes	2 (8.3%)
	No	22 (91.7%)
Allergy	Yes	11 (45.8%)
	No	13 (54.2%)
Gastrointestinal diseases	Yes	19 (79.2%)
	No	5 (20.8%)
Unilateral chewing	Yes	22 (91.7%)
	No	2 (8.3%)
Bruxism	Yes	18 (75%)
	No	6 (25%)
Presence of missing teeth	Yes	13 (54,2%)
	No	2 (45,8%)
Parafunctional habit	Yes	24 (100%)
	No	0 (0%)

X: Mean, SD: Standard Deviation, BMI: Body Mass Index.

Table 2. Pre and post treatment in Orofacial Visual Analogue Scale, all mandibular joint range of motion, pressure pain threshold in masseter, temporalis, and SIPS, Patrick's Faber test, and Oswestry Disability Index

Parameter	X ± SD Median (Min-Max)		P value
	Pre-treatment	Post-treatment	
Orofacial VAS	7.7 ± 1.1	3.7 ± 1.6	0.000 ^a
	8 (6-10)	4 (0-7)	
Depression ROM, cm	4.2 ± 0.4	4.7 ± 0.3	0.000 ^a
	4.2 (3.2-5.2)	4.6 (3.9-5.6)	
Protrusion ROM, cm	0.4 ± 0.2	0.5 ± 0.2	0.000 ^a
	0.4 (0.1-0.8)	0.5 (0.3-1)	
Right lateral deviation ROM, cm	0.7 ± 0.3	0.9 ± 0.2	0.000 ^a
	0.7 (0.1-1.2)	1 (0.5-1.3)	
Left lateral deviation ROM, cm	0.8 ± 0.1	1 ± 0.1	0.000 ^a
	0.8 (0.5-1.1)	1 (0.6-1.4)	
PPT in masseter	4.5 ± 1.4	9.1 ± 3.2	0.000 ^a
	4.5 (2.3-7.5)	8.5 (5-17.1)	
PPT in temporalis	5.3 ± 2.2	9.6 ± 4	0.000 ^a
	4.9 (2.5-10.1)	9.1 (4.5-17.9)	
PPT in SIPS	5.5 ± 2.4	9 ± 2.7	0.000 ^a
	5.3 (1.1-11.4)	8.1 (4.8-15)	
Patrick's faber test, cm	18.1 ± 0.8	15.3 ± 0.9	0.000 ^a
	17.8 (16.9-19.8)	15 (13.8-17.3)	
ODI	39.6 ± 7.3	29.2 ± 5.1	0,000 ^a
	41 (24-52)	29 (18-40)	

a: Paired Samples Test, X: Mean, SD: Standard Deviation, VAS: Visual Analogue Scale, ROM: Range of Motion, PPT: Pressure Pain Threshold, SIPS: Spina Iliaca Posterior Superior, ODI: Oswestry Disability Index, Bold emphasis indicates p<0.05, P significance level.

IBM Statistical Package for Social Sciences 24 (SPSS, Chicago, USA) statistical program was used for data analysis. Continuous variables were given as mean $\pm$ standard deviation, and qualitative variables were given as numbers and percentages (%). It was determined that the data showed normal distribution with the Shapiro–Wilk test, and the relationship between the variables was evaluated with Pearson correlation analysis. Paired Samples test was used to compare the evaluation results obtained before and after treatment. Statistical significance was taken as $p < 0.05$ in all measurements.

Reporting Guidelines

The study was reported in accordance with STROBE guidelines.

Results

Out of the 53 who volunteered to participate in the study, 29 did not meet the inclusion criteria (Figure 2). Twenty-one of the participants were women (87.5%). The median age was 34.8 years and the median BMI was 26.2 kg/m². 41.7% of the participants had a history of trauma, 8.3% had a history of surgery, 45.8% had allergies, 79.2% had gastrointestinal diseases, 91.7% chewed unilaterally, 70% had missing teeth, 100% had parafunctional habit and 75% had bruxism (Table 1). Statistically significant improvement were observed in the Orofacial VAS scores, all mandibular joint ROM values, PPT in masseter, temporalis after treatment in all participants ($p < 0.001$). Also, statistically significant improvement were observed in the PPT in SIPS, Patrick's Faber test score, and ODI score after treatment in all participants ($p < 0.001$), (Table 2). Positive significant relationship were found between the PPT in masseter, temporalis, and SIPS ($r = 0.408, 0.411$; $p = 0.48, 0.46$).

Discussion

The relationship between the craniomandibular system and the SIJ and the mechanisms connecting the two remain unclear. To the best of our knowledge, this study is the first clinical research investigating the effects of OMT applied to the TMJ not only on TMD symptoms but also on SIJ symptoms.

After a single-session treatment, participants showed significant improvements in pain levels, pressure pain thresholds of the masseter and temporalis muscles, and all ranges of motion ROM of the TMJ. Furthermore, enhancements in orofacial pain, all TMJ ROM values, and the pressure pain thresholds of the masseter and temporalis—resulting from orofacial manual therapy—also contributed to better mobility and pressure pain thresholds of the SIPS, along with improved overall functionality.

Although the relationship between TMD and SJD is difficult to explain, a study conducted on healthy individuals⁵ showed that hypomobility functional abnormalities of the SIE emerged after artificial occlusal intervention and suggested that there was a correlation between functional abnormalities of the TMJ and SIJ, which were thought to result from ventrodorsal muscle imbalances. In a study comparing the three-dimensional motion SIEs of patients with TMJ disorders and healthy individuals, it was found that especially joint disc disorders (the most common anterior disc displacement with reduction) may alter the kinematics of the sacroiliac joint.¹⁶ This result indicates that there are statistically significant differences in SIJ kinematics between patients with TMD and healthy individuals. However, it remains unclear whether TMD is the cause or the consequence of these changes. In another study, the function of SIEs was evaluated using

functional tests before and after stabilizing the abnormal position of the mandibular condyle in patients with TMD with the use of a temporary silicone occlusal splint.¹⁷ Significant improvements in test results were observed following the stabilization of the mandibular condyle. Several case studies have also suggested a correlation between TMD and SIJD.^{6–7} Chinappi et al. observed that simultaneous chiropractic and orthodontic treatment in a patient presenting with SIJD complaints resulted in an improvement in SIJD symptoms and indicated a potential connection between these two areas.⁶ In a study where neural therapy was applied to the TMJ, masseter muscle, and C4 level of a patient with both TMD and SIJD, a reduction in the patient's SIJD symptoms was observed.⁷ In this study, improvements in SIJ hypomobility and pressure pain threshold were obtained after OMT was applied to patients with both TMD and SIJD. This results demonstrate a coexistence of these two dysfunctions. However, further research is needed to determine their causative relationships and outcomes.

In the study by Oliveira et al., the effects of occlusal splint and physiotherapy treatment on postural balance were investigated in patients with TMD using a stabilometric platform.¹⁸ A statistically significant increase in antero-posterior velocity was observed in all patients, indicating that this result impacted postural balance. Chung et al., who examined changes in the gait of patients with lower extremity disorders using exercise equipment for TMJ, concluded that the pelvic center point swayed less during the one-legged stance test due to increased stability after treatment.¹⁹ They also reported statistically significant differences in hip, knee, and ankle angles during walking, with these angles returning to the normal range. Tecco et al. found that TMJ disorders were associated with statistically significant changes in walking function based on gait analysis.²⁰ Although gait analysis was not used in this study, a questionnaire addressing daily life functions—such as walking, sitting, and carrying loads—was utilized. Significant improvements were noted in patients' daily life functionality following treatment. These results can reflect the relationship between TMD and SIJD and its impact on functionality.

Manual therapy approaches have an important role in TMJ function, ROM and pain.^{9,21} In this study, after the treatment of orofacial pain, pressure pain threshold in masseter, and temporalis muscles, and ROM values improved in participants. These results may occur due to pain modulation via activating low-threshold A β fibers pathway. This inhibits nociceptive stimuli from A δ and C supply fibers. MT can also elicit affective responses activating opioid, oxytocin, and dopaminergic pathways.²² Due to these effects, several theories exist regarding how improvements in TMD achieved through OMT may impact the SIJ. The relationship between TMJ and SIJ is based on diaphragms, neural connections, and the presence of myofascial chains. The trigeminal nerve is significant due to its role in maintaining postural control, while fascia is of potential importance because it can transmit tension and, through its proprioceptive and nociceptive functions, contribute to referred pain and functional impairments in distant anatomical structures.⁹ Additionally, there are five anatomical diaphragms in the body. These are: the diaphragm muscle, the pelvic floor, the floor of the mouth, the thoracic outlet, and the tentorium of the cerebellum. Previous work has shown the connections between all these diaphragms, with links to fascial and neurological continuity.²³ The function of these diaphragms is to regulate and respond to any changes in intra-abdominal pressure. For instance, if the main inspiratory muscle descends during inhalation, there will be a corresponding lowering of the pelvic floor. With all these connections, symptoms may also manifest in areas far from the source of the problem.⁸

The sacroiliac joint is a gliding joint formed by the iliac and sacral bones, reinforced by the interosseous sacroiliac ligaments, the anterior sacroiliac ligaments, and the dorsal sacroiliac ligaments. The range of motion at the sacroiliac joint is minimal. Therefore, any disorder of this joint and its connections with the structures mentioned above leads to dysfunction.²⁴

Our study has several limitations. It should be recognized that we assessed immediate effects of OMT on orofacial pain, pressure pain threshold, TMJ range of motion, hipomobility in SIJ, and functionality. Immediate changes occurred after OMT treatment provides impetus for future research in this area. Future studies should investigate the long-term effects of this intervention using larger sample sizes in randomized controlled designs. Another limitation is that we cannot rule out the placebo effect due to the absence of a control group. The wide age range of the participants and the lack of a comparison between genders are additional limitations. Future studies should include participants within a specific age range and investigate the effects of gender.

In summary, comprehensive evaluation of patients diagnosed with TMD and considering the patient and their body as a whole is important in the treatment of TMD. Since the body is interconnected kinematically, specific regions shown to have intimate, yet distal, involvement such as the TMJ and pelvis, warrant close clinical attention. With a specific group of patients evaluation and treatment of the TMJ may be essential for a successful clinical outcome in treatment of the sacroiliac joint. Conversely with a subset of some patients, evaluation and treatment of the sacroiliac joint may be essential for a successful clinical outcome in treatment of the stomatognathic dysfunction.

Conclusion

Therefore, when faced with a patient with TMD symptoms, it would be reasonable to question SIJD complaints and include an examination of the TMJ followed by investigations of these anatomical areas where it is deemed appropriate. As the symptoms of TMJ improved, the pressure pain threshold and the mobility of the SIJ also improved.

Clinical Implication

- One session of orofacial manual therapy can improve symptoms of temporomandibular disorder (TMD).
- As pressure pain threshold (PPT) values in the temporomandibular joint (TMJ) improved, PPT values in the sacroiliac posterior superior (SIPS) also improved.
- In patients admitted to the hospital with sacroiliac joint dysfunction (SIJD), it is recommended to evaluate the temporomandibular joint (TMJ) to achieve more effective treatment results.

Declarations

Animal and Human Rights Statement

All procedures performed in this study were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Informed Consent

Written informed consent was obtained from all participants.

Data Availability

The data supporting the findings of this study are available from the corresponding author upon reasonable request.

Conflict of Interest

The authors declare that there is no conflict of interest.

Funding

None.

Abbreviations

- OMT: Orofacial manual therapy
- PPT: Pressure pain threshold
- ROM: Range of motion
- SIJ: Sacroiliac joint
- SIJD: Sacroiliac joint dysfunction
- TMD: Temporomandibular disorder
- TMJ: Temporomandibular joint
- VAS: Visual analog scale

References

1. Kohler AA, Hugoson A, Magnusson T. Prevalence of symptoms indicative of temporomandibular disorders in adults: cross-sectional epidemiological investigations covering two decades. *Acta Odontol Scand.* 2012;70(3):213-223.

2. de Leeuw R, Klasser GD, eds. *Orofacial Pain: Guidelines for Assessment, Diagnosis, and Management.* American Academy of Orofacial Pain. Quintessence Publishing; 2018:122-128. doi:10.25241/stomaeduj.2015.2(2).bookreview.6

3. Ergezen G, Şahin M, Algun ZC. Investigation of the relationship between temporomandibular disorder and postural analysis. *Kocaeli Med J.* 2021;10(2):27-32. doi:10.5505/ktd.2021.22043

4. Kiapour A, Joukar A, Elgafy H, Erbulut DU, Agarwal AK, Goel VK. Biomechanics of the sacroiliac joint: anatomy, function, biomechanics, sexual dimorphism, and causes of pain. *Int J Spine Surg.* 2020;14(1):3-13. doi:10.14444/6077

5. Fink M, Wähling K, Stiesch-Scholz M, Tschernitschek H. The functional relationship between the craniomandibular system, cervical spine, and the sacroiliac joint: a preliminary investigation. *Cranio.* 2003;21(3):202-208. doi:10.1080/08869634.2003.11746252

6. Chinappi AS Jr, Getzoff H. Chiropractic/dental cotreatment of lumbosacral pain with temporomandibular joint involvement. *J Manipulative Physiol Ther.* 1996;19(9):607-612.

7. Senlikci HB, Ozen S. Sacroiliac joint dysfunction treated using neural therapy to the temporomandibular joint: a case report. *Complement Med Res.* 2021;28(4):379-381.

8. Bordoni B, Zanier E. The continuity of the body: hypothesis of treatment of the five diaphragms. *J Altern Complement Med.* 2015;21(4):237-242. doi:10.1089/acm.2013.0211

9. La Touche R, García SM, García BS, et al. Effect of manual therapy and therapeutic exercise applied to the cervical region on pain and pressure pain sensitivity in patients with temporomandibular disorders: a systematic review and meta-analysis. *Pain Med.* 2020;21(10):2373-2384. doi:10.1093/pm/pnaa021

10. Bijur PE, Silver W, Gallagher EJ. Reliability of the visual analog scale for measurement of acute pain. *Acad Emerg Med.* 2001;8(12):1153-1157. doi:10.1111/j.1553-2712.2001.tb01132.x

11. Walker N, Bohannon RW, Cameron D. Discriminant validity of temporomandibular joint range of motion measurements obtained with a ruler. *J Orthop Sports Phys Ther.* 2000;30(8):484-492. doi:10.2519/jospt.2000.30.8.484

12. Stuginski-Barbosa J, Silva RS, Cunha CO, et al. Pressure pain threshold and pain perception in temporomandibular disorder patients: is there any correlation? *Rev Dor.* 2015;16(1):22-26. doi:10.5935/1806-0013.20150005

13. Yakut E, Duger T, Öksüz Ç, et al. Validation of the Turkish version of the Oswestry Disability Index for patients with low back pain. *Spine.* 2004;29(5):581-585. doi:10.1097/01.brs.0000113869.13209.03

14. Tuncer A, Ergun N, Karahan S. Temporomandibular disorders treatment: comparison of home exercise and manual therapy. *Fizyoter Rehabil.* 2013;24(1):9-16. doi:10.1016/j.jbmt.2012.10.006

15. Kalamir A, Pollard H, Vitiello A, Bonello R. Intra-oral myofascial therapy for chronic myogenous temporomandibular disorders: a randomized, controlled pilot study. *J Man Manip Ther.* 2010;18(3):139-146. doi:10.1179/106698110x12640740712374

16. Parchekouhi AB, Karimi N, Barzegar A, Rahnama L. The association between temporomandibular disorders and kinematics of the sacroiliac joint: a 3D motion analysis study. *Elsevier BV.* 2023.

17. Darstka AA, Brzózka M, Bitenc-Jasiejko A, et al. Cause-effect relationships between painful TMD and postural and functional changes in the musculoskeletal system: a preliminary report. *Pain Res Manag.* 2022;2022:1-14.

18. Oliveira SS, Pannuti CM, Paranhos KS, et al. Effect of occlusal splint and therapeutic exercises on postural balance of patients with signs and symptoms of temporomandibular disorder. *Clin Exp Dent Res.* 2019;5(2):109-115. doi:10.1002/cre2.136

19. Chung GY, Choi GS, Shin GY, Park JS. Gait changes after using a temporomandibular joint exerciser in patients who underwent lower limb joint surgery. *J Phys Ther Sci.* 2016;28(5):1584-1587. doi:10.1589/jpts.28.1584

20. Tecco S, Tetè S, D'Attilio SM, Festa F. The analysis of walking in subjects with and without temporomandibular joint disorders: a cross-sectional analysis. *Minerva Stomatol.* 2008;57(9):399-411.

21. Asquini G, Pitance L, Michelotti A, Falla D. Effectiveness of manual therapy applied to craniomandibular structures in temporomandibular disorders: a systematic review. *J Oral Rehabil.* 2022;49(4):442-455. doi:10.1111/joor.13299

22. Armijo-Olivo S, Pitance L, Singh V, Neto F, Thie N, Michelotti A. Effectiveness of manual

therapy and therapeutic exercise for temporomandibular disorders: systematic review and meta-analysis. *Phys Ther.* 2016;96(1):9-25. doi:10.2522/ptj.20140548

23. Bordoni B, Zanier E. The anatomical connections of the diaphragm: the influence of respiration in the body system. *J Multidiscip Healthc.* 2013;6:281-291. doi:10.2147/jmdh.s45443

24. Vleeming A, Schuenke A, Masi T, et al. The sacroiliac joint: an overview of its anatomy, function and potential clinical implications. *J Anat.* 2012;221(6):537-567. doi:10.1111/j.1469-7580.2012.01564.x

How to Cite

Sultan İğrek, Tuğba Kuru Çolak, İlker Çolak, Erdiñ Çekiç. Is there a relationship between temporomandibular disorder and sacroiliac joint dysfunction? *Ann Clin Anal Med* 2025;16(2):99-104. doi:10.4328/ACAM.22373