

Supplementary Table 1. Clinical features of the patient groups

Patient (n = 37%)	
Systemic disease (n = 32 / 37)	
Cardiovascular disease (e.g., CAD, HT, CHF)	17 (%45.9)
Diabetes Mellitus	2 (%5.4)
Chronic renal failure	9 (%24.3)
Inflammatory diseases	2 (%5.4)
Hypo/hyperthyroidism	5 (%13.5)
Triggering Factors (n = 20 / 37)	
Infections	12 (%32.4)
Inflammatory diseases	4 (%10.8)
Drugs	4 (%10.8)
Neoplasms	1 (%2.7)
Surgical procedures	2 (%5.4)
Elementary skin lesions	
Petechiae-purpura	37 (%100)
Urticarial, annular, targetoid papules/plaques	3 (% 8.1)
Vesiculo-bullous lesions	10 (%27.0)
Others (Livedo racemosa, Retiform purpura, Subcutaneous nodules, Digital necrosis, Ulcer)	9 (%24.3)
Site of involvement	
Infra genu	22 (%59.4)
Supra genu-infra umbilical	11 (%29.72)
Supra umbilical	4 (%10.81)
Systemic Symptoms (n = 29 / 37)	
Fever	10 (%27.0)
Fatigue	12 (%32.4)
Arthralgia	24 (64.8)
Myalgia	20 (%54.0)
Weight loss	3 (%8.1)
Abbreviations: CAD = coronary artery disease; HT = hypertension; CHF = congestive heart failure	